



Garlic (*Allium Sativum* Linn.) Extract Supplementation Alters The Free Amino Acid Level In The Gonads Of Male Albino Rats

Sashank Srivastava*, P. H. Pathak

Correspondent Authors:- Sashank Srivastava Department of Zoology, D. D. U. Gorakhpur University, Gorakhpur – 273 009, U. P. India

Email: sashank_sri@rediffmail.com

Effect of Allium sativum Linn. (Alliaceae) (Raw Garlic Extract (RGE)) was studied on free amino acids (FAA) level in male albino rats. The garlic extract was tested in three different doses 1ml, 2ml and 4ml/ kg body weight daily for a period of 7, 14, 21 and 28 days. The significant ($P < 0.01$) increase in Free amino acid (FAA) level was observed when rats were fed with low and medium dose but when rats were fed with 4ml/kg body weight of garlic extract there was a not significant increase in Free Amino Acid (FAA) level was observed.

Key Words: Garlic Extract, Free Amino Acid, Albino Rats, Biochemical Estimation.

INTRODUCTION

Epidemiological studies in the past 10 years have revealed an inverse relationship between *Allium sativum* (garlic) consumption and the incidence of certain diseases, including stomach, colon and laryngeal cancers (Yu-Yan Yeh and Lijuan, 2001). The importance of garlic was recognized many centuries ago, in early Egyptian, Chinese and Indian civilizations as a herbal or traditional medicine. Today, in many parts of the world, garlic is being used for prophylaxis and for the treatment of acute and chronic gastritis, dysentery, typhoid fever, cholera, tuberculosis, pneumonia, diabetes mellitus and heart disease (Khataibeh, *et al.*, 2006). Allicin, diallyldisulfide-oxide, alliin, active ingredients from garlic (alliin) are systemic vasodilator (Sang *et al.*, 1995). Ether extracts and partially purified distilled extracts of garlic have been reported to inhibit human platelet aggregation *in vitro* (Apitz-Castro *et al.*, 1983). Treatment with garlic extracts was found to improve the activation of natural killer cells, the function of T-lymphocytes (Tang *et al.*, 1997). In *in vitro* and *in vivo* studies, aged garlic extracts stimulated immune functions. Kamin and Handler (1951), reported that metabolism of the amino acids follows their accumulation in various tissues. By-products of such metabolism are urea and ammonia which appear in the urine. Bose *et al.*, (1973), reported that glycine labeled with carbon-14 is incorporated into liver cholesterol and fatty acids by rats. Caspi *et al.*, (1971) also reported that rats incorporated the S-methyl group of methionine into cholesterol. Medicinal uses of *Allium sativum* (garlic) have existed for centuries, but its therapeutic properties need more investigation. The present study investigates the effect of garlic extract on amino acid level in both male and female albino rats.

MATERIALS AND METHODS

The Extract

Six months old (after harvest) garlic bulbs were collected from the local market. Garlic bulbs were separated, peeled and washed with distilled water. After drying in shed, about 500 gm of clean garlic bulbs were crushed with the help of electronic grinder. The extract was strained through muslin cloth after squeezing the crushed materials (Sonepati *et al.*, 2001).

Experimental Animal

Healthy adult male albino rats weighing approximately 150 – 200 gm were selected for the experiment. All animals were acclimatized for a week in the laboratory before use (Parthasarathy and Prasanth, 2009). The animals were housed five per cage under controlled conditions of a 12 h light/dark cycle, 50% of humidity and $26^{\circ}\text{C} \pm 2^{\circ}\text{C}$, with minimum noise levels (Nagaraja *et al.*, 2006). Animals had free access to tap water *ad libitum* and normal diet.

Experimental Design

The animals were divided into four groups. Group A animals, which served as healthy control, were given normal feed and tap water *ad libitum* throughout the experimental tannure. Rats of group B, C and D were fed with 1ml, 2ml and 4ml/ kg body weight garlic extract daily for 7, 14, 21 and 28 days daily. In all the groups, the extract was forced fed by using ball – tipped needle every day between 11.00 a.m. to 12.00 pm (Thomson *et al.*, 2006; Cikler *et al.*, 2005).

Free Amino Acids Estimation

Estimation of total free amino acids in gonads of Albino rats was made according to the method of Spies (1957). The tissue homogenate (1% w/v) was prepared in 96% ethanol in an electrical homogenizer for 5 minutes and centrifuged at 8000g for 20 minutes. In 0.1 ml of supernatant, 0.1 ml of distilled water and 2.0 ml of ninhydrin reagent was added and mixed thoroughly. Ninhydrin reagent was prepared by mixing one gram of ninhydrin in 250 ml of absolute ethanol and 0.04 gm of stannous chloride in 25.0 ml of citrate buffer (p^{H} 5.0). The reaction mixture was kept in boiling water bath for 15 minutes. Two ml of 50% ethanol was added to the above after cooling. The violet colour developed which was measured at 575 nm. Glycine was used to determine standard. Free amino acids have been expressed as $\mu\text{g} / \text{mg}$ tissue.

Statistical Analysis

All the experiments were replicated five times and subjected to statistical analysis by two way analysis of variance (ANOVA), followed by student's t-test, wherever required (Paterson, 1939).

RESULTS AND DISCUSSION

There was a significant increase in the mean values of Free Amino Acid level in male albino rats. In group B and C, the level increased significantly (P<0.01) to the extent of 14.81% and 17.42% respectively whereas in group D there was a not significant increase of 1.42% in FAA level was observed

REGIMENS	TREATMENTS	DAYS			
		07	14	21	28
Free Amino Acid level (µg/mg)	CONTROL (0)	304.132 ± 1.282 (100%)	309.207 ± 1.227 (100%)	312.967 ± 1.808 (100%)	317.290 ± 1.282 (100%)
	1ml/kg (bd.wt)	313.531 ± 1.012* (3.09%) ↑	332.703 ± 1.296** (7.60%) ↑	340.974 ± 1.527** (8.95%) ↑	364.282 ± 1.349** (14.81%) ↑
	2ml/kg (bd.wt)	322.929 ± 1.748** (6.18%) ↑	351.124 ± 1.474** (13.56%) ↑	359.771 ± 1.407** (14.95%) ↑	372.553 ± 1.533** (17.42%) ↑
	4ml/kg (bd.wt)	306.688 ± 1.394 ^{NS} (0.74%) ↑	312.027 ± 1.783 ^{NS} (0.91%) ↑	316.350 ± 1.316 ^{NS} (1.08%) ↑	321.801 ± 1.589 ^{NS} (1.42%) ↑

- Values are expressed as Mean ± SE of five replicates, Values in parentheses are percent change with control taken as 100 percent.
- Data were analyzed through Two Way Analysis of Variance (ANOVA) followed by Student’s t-test.
- ‘NS’ not significant, ‘*’ significant (P<0.05) and ‘**’ significant (P<0.01), when treated groups were compared with controls.
- ↓ Decrease in % change and ↑ Increase in % change.




Figure 1: Change in percent level of Free Amino Acid (FAA) in male albino rats after fed with different volumes of raw garlic extract for 7, 14, 21 and 28 days daily.

Amino acids can be metabolized to **produce energy**. Some amino acids act as **neurotransmitters**, and some act as starting materials for the **biosynthesis** of neurotransmitters, hormones, and other important biochemical compounds. Amino acids are the primary building blocks for **proteins**. This description fits a very large number of compounds, most of which are non-physiological. In addition, many physiologically important amino acids are not used in proteins. However, in biochemistry, the term “amino acid” generally refers to one of the 20 types of monomeric units most commonly used to construct proteins. We have no adequate explanation for the changes in amino acid concentrations. A number of possibilities have been considered to explain the mechanism at work but we have as yet no actual proof to substantiate them. This is especially important during fasting, when the breakdown of muscle protein is a major energy source. The increased Amino Acid level suggests possible tissue damage probably due to increased proteolytic activity under stress. However, the elevated levels of amino acid may be **(a)** utilized for energy production by feeding them into the TCA cycle through amino transferase reaction, **(b)** attributed to the synthesis of amino acids in addition to their elevation by protein hydrolysis, **(c)** due to transamination and animation of keto acids, and **(d)** attributed to use of amino acids and their involvements in the maintenance of an acid – base balance.

ACKNOWLEDGEMENT

Authors (PHP) are thankful to University Grant Commission, New Delhi, Letter No. F. No. 34- 414/ 2008 (SR) for their financial support.

REFERENCES

1. Apitz-Castro R, Cabrera S, Cruz MR, Ledezma E and Jain MK. Effects of garlic extracts and of these pure components isolated from it on human platelet aggregation, arachidonate metabolism, release reaction and platelet ultrastructure. *Thromb Res.* (1983); 32: 155.
2. mBose AK, Kapoor NK and Nityanand S. Incorporation of glycine ^{2}C into cholesterol and fatty acids in rats. *Ind. J. Exp. Biol.* (1973); 11: 452-454.
3. Caspi E, Jones JGL and Heidel SP. In vivo incorporation of the S-methyl group of methionine into cholesterol in rats. *Chem. Comm.* (1971); 1201-1202.
4. Çikler E, Saglam B, Zeybek A, Ercan F, Çetinel S, Sener G. The protective effects of aqueous garlic extract against water avoidance stress-induced mast cell degranulation in dermis. *Marmara Medical Journal*, (2005); 18(3): 103-108.

5. Kamin H and Handler P. Metabolism of parenterally administered amino acids. II. Urea synthesis. J. Biol. Chem. (1951); 188: 193-205.
6. Khataibeh M, Abu-Samak M and Banu N. Biochemical investigation of the effect of Garlic (*Allium sativum*) on 7, 12-dimethylbenz[α]anthracene (DMBA) induced mammary cancer in female albino rats. Asian J. Biochemistry (2006); 1(3): 251-256.
7. Nagaraja HS, Anupama BK and Jeganathan PS. Stress response in albino rats. Thai Journal of Physiological Sciences, (2006); 19: 8-15.
8. Parthasarthy G and Prasanth VV. Hepatoprotective Activity of Cassia fistula Linn. Bark extract against Carbon Tetra Chloride Induced Liver Toxicity in Rats, The Internet Journal of Pharmacology, (2009); Vol 6: No. 2.
9. Paterson DD. Statistical Technique in Agricultural Research M.C. Graw- Hill New York (1939).
10. Sang GK, Nam SY, Chung HC, Hongand SY, Jung KH. Enhanced effectiveness of dimethyl-4,4¹-dimethoxy-5,6, 5¹, 6¹,-dimethylene dioxybiphenyl – 2, 2¹ – dicarboxylate in
11. combination with garlic oil against experimental hepatic injury in rats and mice. J. Pharm. Pharmacol. (1995); 47: 678-682.
12. Sonepati SK, Dey S, Dwivedi SK and Swarup D. Effect of garlic (*Allium sativum* L.) extract on tissue lead level in rats. Journal of Ethnopharmacology, (2001); (76): 229 – 232.
13. Spies JR. Colorimetric procedure for amino acids. In “Methods in Enzymology” (eds., SP Colowick and NO Kaplan), Academic Press, (1957); pp 464.
14. Tang ZZ, Sheng S, Liu X, Jian K, Suin and Yan M. Preventing function of garlic on experimental oral precancer and its effect on natural killer cells. Bulletin of Human Medical University. (1997); 22: 312 – 318.
15. Thomson M, Khaled K Al – Qattam, Tanuja Bordia and Muslim Ali. Including Garlic in the Diet May Help Lower Glucose, Cholesterol and Tryglycerides, The Journal of Nutrition, (2006); 136: 800S-802S).
16. Yeh Y and Lijuan L. Cholesterol-Lowering Effect of Garlic Extract and Organosulfur Compounds: Human and Animal Studies, The Journal of Nutrition, (2001); 131: 989S-993S.
17. Table 1: Percent change in Free Amino Acid (FAA) level after following the programmed feeding of *Allium sativum* (garlic) extract daily for 7, 14, 21 and 28 days respectively in male albino rats.
18. Table 1: Percent change in Free Amino Acid (FAA) level after following the programmed feeding of *Allium sativum* (garlic) extract daily for 7, 14, 21 and 28 days respectively in male albino rats.